



CABENUVA
cabotegravir 200 mg/mL; rilpivirine 300 mg/mL
extended-release injectable suspensions

You're on your way.



Welcome to long-acting HIV treatment



CABENUVA

cabotegravir 200 mg/mL; rilpivirine 300 mg/mL
extended-release injectable suspensions

It's time. A different way to treat HIV.

Instead of an HIV treatment you take every day, CABENUVA is an injectable treatment you get just once a month.

With once-a-month CABENUVA, you're good to go.

CABENUVA is given monthly by a healthcare provider as two injections, after you take about a month of once-daily starter pills. It's important to attend all appointments.

Sobek

Undetectable
since 2010



Every person featured in this brochure has been compensated by ViiV Healthcare.

What is CABENUVA?

CABENUVA is a complete prescription regimen used to treat HIV-1 infection in adults as a replacement for their current HIV-1 treatment when their healthcare provider determines that they meet certain requirements.

IMPORTANT SAFETY INFORMATION

Do not receive CABENUVA if you:

- have ever had an allergic reaction to cabotegravir or rilpivirine.
- are taking certain medicines:
 - carbamazepine
 - oxcabazepine
 - phenobarbital
 - phenytoin
 - rifabutin
 - rifampin
 - rifapentine
 - dexamethasone (more than a single-dose treatment), and/or
 - St John's wort (*Hypericum perforatum*)

Please see Important Safety Information throughout and Important Facts About CABENUVA on the last page of this brochure.

How is CABENUVA different?

CABENUVA is an injectable treatment designed to keep you undetectable* for an entire month.

While you're getting your regular monthly injections, **you won't have to take any more pills to treat your HIV.**

CABENUVA is the **first and only long-acting, complete HIV regimen.**

Ask your doctor if CABENUVA is right for you.

Raphael
Undetectable
since 2009

*Undetectable means the amount of HIV in the blood is below the level that can be measured by a lab test. Results may vary.

IMPORTANT SAFETY INFORMATION (cont'd)

What are the possible side effects of CABENUVA?

CABENUVA may cause serious side effects, including:

- **Allergic reactions.** Call your healthcare provider right away if you develop a rash with CABENUVA. **Stop receiving CABENUVA and get medical help right away if you develop a rash with any of the following signs or symptoms:**
 - fever
 - generally ill feeling
 - tiredness
 - muscle or joint aches
 - trouble breathing
 - blisters or sores in mouth
 - blisters
 - redness or swelling of the eyes
 - swelling of the mouth, face, lips, or tongue
- **Post-injection reactions** have happened within minutes in some people after receiving their rilpivirine injection. Most symptoms resolved within a few minutes after the injection. Symptoms may include:
 - trouble breathing
 - stomach cramps
 - sweating
 - numbness of your mouth
 - feeling anxious
 - feeling warm
 - feeling light-headed or feeling like you are going to pass out (faint)
 - blood pressure changes

Please see Important Safety Information throughout and Important Facts About CABENUVA on the last page of this brochure.

“In my experience, I can get my injections during my lunch break—my doctor gets me in and out most days.”

Mark
Stays undetectable with CABENUVA
Results may vary.



IMPORTANT SAFETY INFORMATION (cont'd)

What are the possible side effects of CABENUVA? (cont'd)

CABENUVA may cause serious side effects, including: (cont'd)

- **Liver problems.** People with a history of hepatitis B or C virus or people who have certain liver function test changes may have an increased risk of developing new or worsening changes in certain liver tests during CABENUVA treatment. Liver problems have also happened in people without history of liver problems or other risk factors. Your healthcare provider may do blood tests to check your liver function. **Call your healthcare provider right away if you develop any of the following signs or symptoms of liver problems:**
 - your skin or the white part of your eyes turns yellow (jaundice)
 - dark or “tea-colored” urine
 - light-colored stools (bowel movements)
 - nausea or vomiting
 - loss of appetite
 - pain or tenderness on the right side of your stomach area
 - itching
- **Depression or mood changes. Call your healthcare provider or get emergency medical help right away if you have any of the following symptoms:**
 - feeling sad or hopeless
 - feeling anxious or restless
 - have thoughts of hurting yourself (suicide) or have tried to hurt yourself

Please see Important Safety Information throughout and Important Facts About CABENUVA on the last page of this brochure.

Scheduling your monthly treatment

Target Treatment Date


You and your doctor will choose one day a month to get your injections. That's called your **Target Treatment Date**. It's important to meet this date each month.

If you can't make your Target Treatment Date, make sure to contact your doctor right away.

Flexible Treatment Window

You have a **Flexible Treatment Window**—from 7 days before to 7 days after your Target Treatment Date.

To help stay undetectable, it's important to keep your monthly appointments and stick with your treatment plan.



- target treatment date
- flexible treatment window

Please see Important Safety Information throughout and Important Facts About CABENUVA on the last page of this brochure.

IMPORTANT SAFETY INFORMATION (cont'd)

What are the possible side effects of CABENUVA? (cont'd)

The most common side effects of CABENUVA include:

- pain, tenderness, hardened mass or lump, swelling, redness, itching, bruising, and warmth at the injection site
- fever
- tiredness
- headache
- muscle or bone pain
- nausea
- sleep problems
- dizziness
- rash

These are not all the possible side effects of CABENUVA. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Michelle
Undetectable
since 1998



What the clinical studies show

Once-a-month CABENUVA is proven to be as effective as daily HIV pills at keeping people undetectable.*†

CABENUVA was tested
in 2 clinical studies:

With more than
1,000 adults

Over a
48-week period

People were already
undetectable and divided
into 2 equal groups:

Those who switched
to CABENUVA

Those who continued on their
current daily pill regimen*

The studies included a **diverse range of people** across age, race, and gender, including transgender individuals.

At Week 48:

- **9 out of 10 people** remained undetectable, whether they were on daily HIV pills or CABENUVA
- **Less than 2% of people** did not remain undetectable [primary endpoint]

*Daily pill regimens tested contained dolutegravir and 2 nucleoside reverse transcriptase inhibitors (NRTIs) or 2 NRTIs plus a protease inhibitor (PI), NNRTI (non-nucleoside reverse transcriptase inhibitor) or INSTI (integrase strand inhibitor).

†Undetectable means the amount of HIV in the blood is below the level that can be measured by a lab test (less than 50 copies/mL).

*Results may vary.

IMPORTANT SAFETY INFORMATION (cont'd)

Before you receive CABENUVA, tell your healthcare provider about all your medical conditions, including if you:

- have ever had a skin rash or an allergic reaction to medicines that contain cabotegravir or rilpivirine.
- have or have had liver problems, including hepatitis B or C infection.
- have ever had mental health problems.
- are pregnant or plan to become pregnant. It is not known if CABENUVA will harm your unborn baby. CABENUVA can remain in your body for up to 12 months or longer after the last injection.
- are breastfeeding or plan to breastfeed. **Do not breastfeed if you take CABENUVA.** You should not breastfeed if you have HIV-1 because of the risk of passing HIV-1 to your baby. It is not known if CABENUVA can pass to your baby in your breast milk. Talk with your healthcare provider about the best way to feed your baby during treatment.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

- Some medicines interact with CABENUVA. Keep a list of your medicines and show it to your healthcare provider and pharmacist when you get a new medicine. You can ask your healthcare provider or pharmacist for a list of medicines that interact with CABENUVA.
- **Do not start taking a new medicine without telling your healthcare provider.** Your healthcare provider can tell you if it is safe to take CABENUVA with other medicines.

Please see Important Safety Information throughout and Important Facts About CABENUVA on the last page of this brochure.

Side effects

The most common side effects reported in clinical studies were injection-site reactions.*

83% of people in the clinical studies experienced injection-site reactions.

99% of injection-site reactions were mild to moderate.

4% of people on CABENUVA stopped treatment due to any side effect.

Aside from injection-site reactions, the most common side effects were:

Side effect	Switched to CABENUVA	Continued on daily pill regimen†
Fever	8%	0%
Tiredness	5%	<1%
Headache	4%	<1%
Muscle or bone pains	3%	<1%
Nausea	3%	1%
Sleep problems	2%	<1%
Dizziness	2%	<1%
Rash	2%	0%

Results may vary.

*Injection-site reactions include pain, tenderness, hardened mass or lump, swelling, redness, itching, bruising, and warmth at the injection site.

†Daily pill regimens tested contained dolutegravir and 2 nucleoside reverse transcriptase inhibitors (NRTIs) or 2 NRTIs plus a protease inhibitor (PI), NNRTI (non-nucleoside reverse transcriptase inhibitor) or INSTI (integrase strand inhibitor).

IMPORTANT SAFETY INFORMATION (cont'd)

How will I receive CABENUVA?

- Once a month, a healthcare provider will inject CABENUVA as 2 injections (cabotegravir and rilpivirine), one into each side of your buttocks.
- Before receiving your first CABENUVA injections, your healthcare provider will have you take one VOCABRIA (cabotegravir) tablet and one EDURANT (rilpivirine) tablet once a day for one month (at least 28 days) to assess how well you tolerate these medicines.
- Stay under the care of a healthcare provider during treatment with CABENUVA. It is important that you attend your planned appointments to receive your injections.
- If you miss or plan to miss a scheduled monthly CABENUVA injection by more than 7 days, call your healthcare provider right away to discuss your treatment options.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Please read the Patient Information for CABENUVA and discuss it with your healthcare provider.

Please see Important Safety Information throughout and Important Facts About CABENUVA on the last page of this brochure.

What people said about switching to CABENUVA



9 out of 10 people who switched to CABENUVA preferred once-a-month injections over their previous daily pill regimen.*†

In a survey conducted during clinical studies, people were asked how they felt about their treatment, including which one they preferred. The results are their opinions and don't imply clinical effectiveness.

"Now that I'm on CABENUVA, I don't have to think about taking my HIV pill every day."

Jakub
Stays undetectable
with CABENUVA



*Daily pill regimens tested contained dolutegravir and 2 nucleoside reverse transcriptase inhibitors (NRTIs) or 2 NRTIs plus a protease inhibitor (PI), NNRTI (non-nucleoside reverse transcriptase inhibitor) or INSTI (integrase strand inhibitor).

†88% of people said they preferred CABENUVA, 2% preferred previous daily regimen, and 10% did not respond to the survey question. Results may vary.

IMPORTANT SAFETY INFORMATION

Do not receive CABENUVA if you:

- have ever had an allergic reaction to cabotegravir or rilpivirine.
- are taking certain medicines:
 - carbamazepine
 - oxcarbazepine
 - phenobarbital
 - phenytoin
 - rifabutin
 - rifampin
 - rifapentine
 - dexamethasone (more than a single-dose treatment), and/or
 - St John's wort (*Hypericum perforatum*)

What are the possible side effects of CABENUVA?

CABENUVA may cause serious side effects, including:

- **Allergic reactions.** Call your healthcare provider right away if you develop a rash with CABENUVA. **Stop receiving CABENUVA and get medical help right away if you develop a rash with any of the following signs or symptoms:**
 - fever
 - generally ill feeling
 - tiredness
 - muscle or joint aches
 - trouble breathing
 - blisters or sores in mouth
 - blisters
 - redness or swelling of the eyes
 - swelling of the mouth, face, lips, or tongue

Please see Important Safety Information throughout and Important Facts About CABENUVA on the last page of this brochure.

Have the CABENUVA conversation with your doctor today

Nobody knows you like you know you. Tell your doctor about your treatment priorities. Together, you can best evaluate whether CABENUVA is a good fit for you.

Here are some topics that can help you and your doctor have a productive discussion about CABENUVA:

How you feel about taking daily HIV pills

What switching to a once-monthly treatment would mean for you

If you're undetectable,* how you can make the switch to CABENUVA

Daily HIV treatment has helped so many control their HIV. Today, you have a treatment option you've never had before: once-monthly CABENUVA.

To learn more about CABENUVA, talk with your doctor.

IMPORTANT SAFETY INFORMATION (cont'd)

What are the possible side effects of CABENUVA? (cont'd)

CABENUVA may cause serious side effects, including: (cont'd)

- **Post-injection reactions** have happened within minutes in some people after receiving their rilpivirine injection. Most symptoms resolved within a few minutes after the injection. Symptoms may include:
 - trouble breathing
 - stomach cramps
 - sweating
 - numbness of your mouth
 - feeling warm
 - feeling light-headed or feeling like you are going to pass out (faint)
 - blood pressure changes
 - feeling anxious
- **Liver problems.** People with a history of hepatitis B or C virus or people who have certain liver function test changes may have an increased risk of developing new or worsening changes in certain liver tests during CABENUVA treatment. Liver problems have also happened in people without history of liver problems or other risk factors. Your healthcare provider may do blood tests to check your liver function. **Call your healthcare provider right away if you develop any of the following signs or symptoms of liver problems:**
 - your skin or the white part of your eyes turns yellow (jaundice)
 - dark or "tea-colored" urine
 - light-colored stools (bowel movements)
 - nausea or vomiting
 - loss of appetite
 - pain or tenderness on the right side of your stomach area
 - itching

Please see Important Safety Information throughout and Important Facts About CABENUVA on the last page of this brochure.

*Undetectable means the amount of HIV in the blood is below the level that can be measured by a lab test.

CABENUVA cost support

ViiVConnect can help you explore ways to pay for your prescribed ViiV Healthcare medications, whether you're insured or uninsured.

If you're prescribed CABENUVA, you can enroll in ViiVConnect to:

- Determine your insurance coverage
- Find out if you're eligible for programs that may help lower your out-of-pocket costs^a



ViiVConnect.com/injectable

Orlando
Stays undetectable
with CABENUVA
Results may vary.

IMPORTANT SAFETY INFORMATION (cont'd)

What are the possible side effects of CABENUVA? (cont'd)

CABENUVA may cause serious side effects, including: (cont'd)

- **Depression or mood changes.** Call your healthcare provider or get emergency medical help right away if you have any of the following symptoms:
 - feeling sad or hopeless
 - feeling anxious or restless
 - have thoughts of hurting yourself (suicide) or have tried to hurt yourself



Check out ways to access your prescribed ViiV Healthcare medications

| Insurance Review

| Financial Assistance Programs^a

Call to speak to an Access Coordinator

Get support from ViiVConnect




1-844-588-3288 (toll-free)
Monday-Friday, 8AM-11PM (ET)

LEARN MORE

^aSubject to eligibility and program terms and conditions;
ViiVConnect programs do not constitute health insurance.

Please see Important Safety Information throughout and Important Facts About CABENUVA on the last page of this brochure.



***Talk with your doctor about
CABENUVA today.***



CABENUVA.com

Important Facts About CABENUVA

This is only a brief summary of important information about CABENUVA and does not replace talking to your healthcare provider about your condition and treatment.

(kab' en ue vah)

ABOUT CABENUVA

CABENUVA is a complete prescription regimen used to treat HIV-1 infection in adults as a replacement for their current HIV-1 treatment when their healthcare provider determines that they meet certain requirements.

HIV-1 is the virus that causes Acquired Immune Deficiency Syndrome (AIDS).

CABENUVA contains 2 different medicines:

- cabotegravir
- rilpivirine

It is not known if CABENUVA is safe and effective in children.

DO NOT RECEIVE CABENUVA IF YOU

- have ever had an allergic reaction to cabotegravir or rilpivirine.
- are taking certain medicines:
 - carbamazepine
 - oxcarbazepine
 - phenobarbital
 - phenytoin
 - rifabutin
 - rifampin
 - rifapentine
 - dexamethasone (more than a single-dose treatment), and/or
 - St John's wort (*Hypericum perforatum*)

BEFORE RECEIVING CABENUVA

Tell your healthcare provider about all your medical conditions, including if you:

- have ever had a skin rash or an allergic reaction to medicines that contain cabotegravir or rilpivirine.
- have or have had liver problems, including hepatitis B or C infection.
- have ever had mental health problems.
- are pregnant or plan to become pregnant. It is not known if CABENUVA will harm your unborn baby. CABENUVA can remain in your body for up to 12 months or longer after the last injection.
- are breastfeeding or plan to breastfeed. **Do not breastfeed if you take CABENUVA.**
 - You should not breastfeed if you have HIV-1 because of the risk of passing HIV-1 to your baby. It is not known if CABENUVA can pass to your baby in your breast milk.
 - Talk with your healthcare provider about the best way to feed your baby during treatment with CABENUVA.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Some medicines interact with CABENUVA. Keep a list of your medicines and show it to your healthcare provider and pharmacist when you get a new medicine. You can ask your healthcare provider or pharmacist for a list of medicines that interact with CABENUVA.

Do not start taking a new medicine without telling your healthcare provider. Your healthcare provider can tell you if it is safe to take CABENUVA with other medicines.

POSSIBLE SIDE EFFECTS OF CABENUVA

CABENUVA may cause serious side effects, including:

- **Allergic reactions.** Call your healthcare provider right away if you develop a rash with CABENUVA. **Stop receiving CABENUVA and get medical help right away if you develop a rash with any of the following signs or symptoms:**
 - fever
 - generally ill feeling
 - tiredness
 - muscle or joint aches
 - trouble breathing
 - blisters or sores in mouth
 - blisters
 - redness or swelling of the eyes
 - swelling of the mouth, face, lips, or tongue
- **Post-injection reactions** have happened within minutes in some people after receiving their rilpivirine injection. Most symptoms resolved within a few minutes after the injection. Symptoms may include:
 - trouble breathing
 - stomach cramps
 - sweating
 - numbness of your mouth
 - feeling warm
 - feeling light-headed or feeling like you are going to pass out (faint)
 - blood pressure changes
- **Liver problems.** People with a history of hepatitis B or C virus or people who have certain liver function test changes may have an increased risk of developing new or worsening changes in certain liver tests during CABENUVA treatment. Liver problems have also happened in people without history of liver problems or other risk factors. Your healthcare provider may do blood tests to check your liver function. **Call your healthcare provider right away if you develop any of the following signs or symptoms of liver problems:**
 - your skin or the white part of your eyes turns yellow (jaundice)
 - dark or "tea-colored" urine
 - light-colored stools (bowel movements)
 - nausea or vomiting
 - loss of appetite
 - pain or tenderness on the right side of your stomach area
 - itching
- **Depression or mood changes.** Call your healthcare provider or get emergency medical help right away if you have any of the following symptoms:
 - feeling sad or hopeless
 - feeling anxious or restless
 - have thoughts of hurting yourself (suicide) or have tried to hurt yourself

The most common side effects of CABENUVA include:

- pain, tenderness, hardened mass or lump, swelling, redness, itching, bruising, and warmth at the injection site
- fever
- tiredness
- headache
- muscle or bone pain
- nausea
- sleep problems
- dizziness
- rash

These are not all the possible side effects of CABENUVA. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

GET MORE INFORMATION

- Talk to your healthcare provider or pharmacist.
- Go to CABENUVA.com or call 1-877-844-8872, where you can also get FDA-approved labeling.

January 2021 CBN:1PIL

Trademark is owned by or licensed to the ViiV Healthcare group of companies.

©2021 ViiV Healthcare or licensor.

CBROGM210040 June 2021

Produced in USA.