



CABENUVA

cabotegravir 200 mg/mL; rilpivirine 300 mg/mL
extended-release injectable suspensions

Heard about injectable CABENUVA?

Find out what you need to know inside.



Every other month, and you're good to go.

CABENUVA is given every other month or monthly by a healthcare provider as two injections, after you take about a month of once-daily starter pills. Your every other month regimen begins after two consecutive months of injections. It's important to attend all appointments.

A different way to treat HIV.



Unlike daily pills, CABENUVA is a long-acting, complete HIV regimen you can get every other month.

Steve

Hiker
Avid traveler
Stays undetectable
with CABENUVA

Results may vary.



People featured are living with HIV and have been compensated by ViiV Healthcare.

What is CABENUVA?

CABENUVA is a complete prescription regimen used to treat HIV-1 infection in adults as a replacement for their current HIV-1 treatment when their healthcare provider determines that they meet certain requirements.

IMPORTANT SAFETY INFORMATION

Do not receive CABENUVA if you:

- have ever had an allergic reaction to cabotegravir or rilpivirine.
- are taking certain medicines: carbamazepine; oxcarbazepine; phenobarbital; phenytoin; rifabutin; rifampin; rifapentine; dexamethasone (more than a single-dose treatment); St John's wort (*Hypericum perforatum*).

Please see Important Safety Information throughout and Important Facts About CABENUVA on the last page of this brochure.

How is CABENUVA different?

CABENUVA is the only complete, long-acting injectable HIV treatment **proven to help keep people undetectable* for up to 2 months**, depending on the treatment plan.

With regular CABENUVA injections, **you won't have to take any more daily HIV pills.**

"Instead of taking my HIV pill every day, I get CABENUVA just once every other month. That's only 6 times a year."

Jayson
Veteran
History buff
Stays undetectable
with CABENUVA
Results may vary.

*Undetectable means the amount of HIV in the blood is below the level that can be measured by a lab test. Results may vary.

IMPORTANT SAFETY INFORMATION (cont'd)

What are the possible side effects of CABENUVA?

CABENUVA may cause serious side effects, including:

- **Allergic reactions.** Call your healthcare provider right away if you develop a rash with CABENUVA. **Stop receiving CABENUVA and get medical help right away if you develop a rash with any of the following signs or symptoms:** fever; generally ill feeling; tiredness; muscle or joint aches; trouble breathing; blisters or sores in mouth; blisters; redness or swelling of the eyes; swelling of the mouth, face, lips, or tongue.
- **Post-injection reactions** have happened within minutes in some people after receiving their rilpivirine injection. Most symptoms resolved within minutes after the injection. Symptoms may include: trouble breathing; narrowing of airways; stomach cramps; sweating; numbness of your mouth; pain (eg, back and chest); feeling anxious; feeling warm; rash; feeling lightheaded or like you are going to faint; blood pressure changes.

Please see Important Safety Information throughout and Important Facts About CABENUVA on the last page of this brochure.

***“I work two jobs,
so long-acting treatment
really works with
my lifestyle.”***

Karen

Lab technician

Proud mom

Stays undetectable
with CABENUVA

Results may vary.



IMPORTANT SAFETY INFORMATION (cont'd)

**What are the possible side effects of CABENUVA?
(cont'd)**

CABENUVA may cause serious side effects, including: (cont'd)

- **Liver problems.** People with a history of hepatitis B or C virus or people who have certain liver function test changes may have an increased risk of developing new or worsening changes in certain liver tests during CABENUVA treatment. Liver problems have also happened in people without history of liver problems or other risk factors. Your healthcare provider may do blood tests to check your liver function. **Call your healthcare provider right away if you develop any of the following signs or symptoms of liver problems:** your skin or the white part of your eyes turns yellow (jaundice); dark or “tea-colored” urine; light-colored stools (bowel movements); nausea or vomiting; loss of appetite; pain, aching, or tenderness on the right side of your stomach area; itching.
- **Depression or mood changes.** **Call your healthcare provider or get emergency medical help right away if you have any of the following symptoms:** feeling sad or hopeless; feeling anxious or restless; have thoughts of hurting yourself (suicide) or have tried to hurt yourself.

**Please see Important Safety Information throughout
and Important Facts About CABENUVA on the last page
of this brochure.**

Scheduling your CABENUVA treatments

CABENUVA injections can be given once a month or every other month, depending on your treatment plan.


To help stay undetectable, it's important to keep your planned appointments and stick to your treatment plan.



Target Treatment Date

You and your doctor will choose an ongoing date that works best to get your injections from a healthcare professional. This is called your **Target Treatment Date**. If you can't make your appointment, be sure to contact your doctor right away.

Flexible Treatment Window

You have a **Flexible Treatment Window**—from 7 days before to 7 days after your Target Treatment Date.



-  Target Treatment Date
-  Flexible Treatment Window

IMPORTANT SAFETY INFORMATION (cont'd)

What are the possible side effects of CABENUVA? (cont'd)

The most common side effects of CABENUVA include:

- Pain, tenderness, hardened mass or lump, swelling, redness, itching, bruising, and warmth at the injection site; fever; tiredness; headache; muscle or bone pain; nausea; sleep problems; dizziness; rash.

These are not all the possible side effects of CABENUVA.

Call your doctor for medical advice about side effects.

You may report side effects to FDA at 1-800-FDA-1088.

Please see Important Safety Information throughout and Important Facts About CABENUVA on the last page of this brochure.

What the clinical studies show

Long-acting CABENUVA is proven to help keep people undetectable.*

CABENUVA was tested in 3 clinical studies, involving more than 2,000 undetectable adults who either switched to CABENUVA or continued their HIV regimen.

In two 48-week studies

About 1,000 people received once-monthly CABENUVA or continued their daily pill regimen.[†]

In another 48-week study

About 1,000 people received CABENUVA injections once a month or every other month.

All studies included a **diverse range of participants** across age, race, and gender, including transgender people.

At Week 48 in all studies:

- **9 out of 10 people** remained undetectable, whether they were on daily HIV pills or CABENUVA
- **Less than 2% of people** did not remain undetectable [primary endpoint]

IMPORTANT SAFETY INFORMATION (cont'd)

Before you receive CABENUVA, tell your healthcare provider about all your medical conditions, including if you:

- have ever had a skin rash or an allergic reaction to medicines that contain cabotegravir or rilpivirine.
- have ever had liver problems, including hepatitis B or C infection.
- have ever had mental health problems.
- are pregnant or plan to become pregnant. It is not known if CABENUVA will harm your unborn baby. CABENUVA can remain in your body for up to 12 months or longer after the last injection.
- are breastfeeding or plan to breastfeed. **Do not breastfeed if you take CABENUVA.** You should not breastfeed if you have HIV-1 because of the risk of passing HIV-1 to your baby. It is not known if CABENUVA can pass to your baby in your breast milk. Talk with your healthcare provider about the best way to feed your baby during treatment.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.


- Some medicines interact with CABENUVA. Keep a list of your medicines and show it to your healthcare provider and pharmacist. **Do not start taking a new medicine without telling your healthcare provider.**

Please see Important Safety Information throughout and Important Facts About CABENUVA on the last page of this brochure.

*Undetectable means the amount of HIV in the blood is below the level that can be measured by a lab test (less than 50 copies/mL). Results may vary.

[†]Daily pill regimens contained dolutegravir and 2 nucleoside reverse transcriptase inhibitors (NRTIs) or 2 NRTIs plus a protease inhibitor (PI), non-nucleoside reverse transcriptase inhibitor (NNRTI), or integrase strand transfer inhibitor (INSTI).

Side effects



The most common side effects reported in all clinical studies were injection-site reactions.*

- 75% to 83% of patients experienced injection-site reactions.

Most injection-site reactions were mild to moderate.

Aside from injection-site reactions, the most common side effects to occur in less than 9% of people were:

- Fever
- Tiredness
- Headache
- Muscle or bone pain
- Nausea
- Sleep problems
- Dizziness
- Rash

Fewer than 4% of people on CABENUVA stopped treatment due to any side effect.

Results may vary.

*Injection-site reactions include pain, tenderness, hardened mass or lump, swelling, redness, itching, bruising, and warmth at the injection site.

IMPORTANT SAFETY INFORMATION (cont'd)

How will I receive CABENUVA?

- Once a month or once every 2 months, a healthcare provider will inject CABENUVA as 2 injections (cabotegravir and rilpivirine), one into the muscle of each side of your buttocks.
- Before receiving your first CABENUVA injections, your healthcare provider will have you take one VOCABRIA (cabotegravir) tablet and one EDURANT (rilpivirine) tablet once a day for 1 month (at least 28 days) to assess how well you tolerate these medicines.
- Stay under the care of a healthcare provider during CABENUVA treatment. It is important that you attend all planned injection appointments.
- If you miss or plan to miss a scheduled CABENUVA injection by more than 7 days, call your healthcare provider right away to discuss your treatment options.
- If you stop treatment with CABENUVA you will need to take other medicines to treat your HIV-1 infection and reduce the risk of developing viral resistance. Call your healthcare provider right away to discuss your treatment options.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Please read the Patient Information for CABENUVA and discuss it with your healthcare provider.

Please see Important Safety Information throughout and Important Facts About CABENUVA on the last page of this brochure.

What people said about switching to CABENUVA



9 out of 10 people preferred the long-acting CABENUVA regimen they switched to over their previous regimen.**

In a survey conducted during clinical studies, people were asked how they felt about their treatment, including which one they preferred. The results are their opinions and don't imply clinical effectiveness.

"HIV pills aren't on my mind anymore."

Orlando
Fashionista
Event planner
Stays undetectable
with CABENUVA

Results may vary.

*88% of people said they preferred once-monthly CABENUVA over their previous daily HIV pills, 2% preferred daily HIV pills, and 10% did not respond to the survey question.

*94% of people with prior CABENUVA experience said they preferred injections every other month over their previous once-monthly injections or daily starter pills, 3% preferred once-monthly injections, 2% preferred daily starter pills, and 1% reported no preference.

*98% of people with no previous CABENUVA use preferred injections every other month over daily starter pills, 1% preferred daily starter pills, and <1% reported no preference.

IMPORTANT SAFETY INFORMATION

Do not receive CABENUVA if you:

- have ever had an allergic reaction to cabotegravir or rilpivirine.
- are taking certain medicines: carbamazepine; oxcarbazepine; phenobarbital; phenytoin; rifabutin; rifampin; rifapentine; dexamethasone (more than a single-dose treatment); St John's wort (*Hypericum perforatum*).

What are the possible side effects of CABENUVA?

CABENUVA may cause serious side effects, including:

- **Allergic reactions.** Call your healthcare provider right away if you develop a rash with CABENUVA. **Stop receiving CABENUVA and get medical help right away if you develop a rash with any of the following signs or symptoms:** fever; generally ill feeling; tiredness; muscle or joint aches; trouble breathing; blisters or sores in mouth; blisters; redness or swelling of the eyes; swelling of the mouth, face, lips, or tongue.
- **Post-injection reactions** have happened within minutes in some people after receiving their rilpivirine injection. Most symptoms resolved within minutes after the injection. Symptoms may include: trouble breathing; narrowing of airways; stomach cramps; sweating; numbness of your mouth; pain (eg, back and chest); feeling anxious; feeling warm; rash; feeling lightheaded or like you are going to faint; blood pressure changes.

Please see Important Safety Information throughout and Important Facts About CABENUVA on the last page of this brochure.

Nobody knows you like you know you

By discussing your treatment priorities with your doctor, you can find out if CABENUVA is right for you.

Have the CABENUVA conversation with your doctor today.

Here are some topics to help get you started:

How you feel about taking daily HIV pills

What switching to a long-acting treatment would mean for you

If you're undetectable,* how you can make the switch to CABENUVA

IMPORTANT SAFETY INFORMATION (cont'd)

What are the possible side effects of CABENUVA? (cont'd)

CABENUVA may cause serious side effects, including: (cont'd)

- **Liver problems.** People with a history of hepatitis B or C virus or people who have certain liver function test changes may have an increased risk of developing new or worsening changes in certain liver tests during CABENUVA treatment. Liver problems have also happened in people without history of liver problems or other risk factors. Your healthcare provider may do blood tests to check your liver function. **Call your healthcare provider right away if you develop any of the following signs or symptoms of liver problems:** your skin or the white part of your eyes turns yellow (jaundice); dark or "tea-colored" urine; light-colored stools (bowel movements); nausea or vomiting; loss of appetite; pain, aching, or tenderness on the right side of your stomach area; itching.

Please see Important Safety Information throughout and Important Facts About CABENUVA on the last page of this brochure.

*Undetectable means the amount of HIV in the blood is below the level that can be measured by a lab test.

Help paying for CABENUVA

ViiVConnect can help you explore ways to pay for your prescribed ViiV Healthcare medications, **whether you're insured or not.**

If you're prescribed CABENUVA, you can enroll in ViiVConnect to:

- Get help navigating the insurance process
- Find out if you're eligible for the CABENUVA Savings Program that may help lower your out-of-pocket costs. If approved, your co-pay could be as little as \$0.^a



ViiVConnect.com

Mark

Nature enthusiast
Adventurer
Stays undetectable
with CABENUVA

Results may vary.



IMPORTANT SAFETY INFORMATION (cont'd)

What are the possible side effects of CABENUVA? (cont'd)

CABENUVA may cause serious side effects, including: (cont'd)

- Depression or mood changes. Call your healthcare provider or get emergency medical help right away if you have any of the following symptoms: feeling sad or hopeless; feeling anxious or restless; have thoughts of hurting yourself (suicide) or have tried to hurt yourself.



Check out ways to access your prescribed ViiV Healthcare medications

Insurance Review

Financial Assistance Programs^a

Call to speak to an Access Coordinator

Get support from ViiVConnect



1-844-588-3288 (toll-free)
Monday-Friday, 8AM-11PM (ET)

LEARN MORE

^aSubject to eligibility and program terms and conditions;
ViiVConnect programs do not constitute health insurance.

CABENUVA and VOCABRIA are trademarks owned by or licensed to the ViiV Healthcare group of companies. The other brand listed is a trademark owned by or licensed to its respective owner, who is not affiliated with the ViiV Healthcare group of companies.

Please see Important Safety Information throughout and Important Facts About CABENUVA on the last page of this brochure.

***Ask your doctor today if
CABENUVA is right for you.***



Visit CABENUVA.com

Important Facts About CABENUVA

This is only a brief summary of important information about CABENUVA and does not replace talking to your healthcare provider about your condition and treatment.

(kab' en ue vah)

ABOUT CABENUVA

CABENUVA is a complete prescription regimen used to treat HIV-1 infection in adults as a replacement for their current HIV-1 treatment when their healthcare provider determines that they meet certain requirements.

HIV-1 is the virus that causes Acquired Immune Deficiency Syndrome (AIDS).

CABENUVA contains 2 different medicines:

- cabotegravir
- rilpivirine

It is not known if CABENUVA is safe and effective in children.

DO NOT RECEIVE CABENUVA IF YOU

- have ever had an allergic reaction to cabotegravir or rilpivirine.
- are taking certain medicines:
 - carbamazepine
 - oxcarbazepine
 - phenobarbital
 - phenytoin
 - rifabutin
 - rifampin
 - rifapentine
 - dexamethasone (more than a single-dose treatment), and/or
 - St John's wort (*Hypericum perforatum*)

BEFORE RECEIVING CABENUVA

Tell your healthcare provider about all your medical conditions, including if you:

- have ever had a skin rash or an allergic reaction to medicines that contain cabotegravir or rilpivirine.
- have ever had liver problems, including hepatitis B or C infection.
- have ever had mental health problems.
- are pregnant or plan to become pregnant. It is not known if CABENUVA will harm your unborn baby. CABENUVA can remain in your body for up to 12 months or longer after the last injection.
- are breastfeeding or plan to breastfeed. **Do not breastfeed if you take CABENUVA.**
 - You should not breastfeed if you have HIV-1 because of the risk of passing HIV-1 to your baby. It is not known if CABENUVA can pass to your baby in your breast milk.
 - Talk with your healthcare provider about the best way to feed your baby during treatment with CABENUVA.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Some medicines interact with CABENUVA. Keep a list of your medicines and show it to your healthcare provider and pharmacist when you get a new medicine. You can ask your healthcare provider or pharmacist for a list of medicines that interact with CABENUVA.

Do not start taking a new medicine without telling your healthcare provider. Your healthcare provider can tell you if it is safe to take CABENUVA with other medicines.

POSSIBLE SIDE EFFECTS OF CABENUVA

CABENUVA may cause serious side effects, including:

- **Allergic reactions.** Call your healthcare provider right away if you develop a rash with CABENUVA. **Stop receiving CABENUVA and get medical help right away if you develop a rash with any of the following signs or symptoms:**
 - fever
 - generally ill feeling
 - tiredness
 - muscle or joint aches
 - trouble breathing
 - blisters or sores in mouth
 - blisters
 - redness or swelling of the eyes
 - swelling of the mouth, face, lips, or tongue
- **Post-injection reactions** have happened within minutes in some people after receiving their rilpivirine injection. Most symptoms resolved within minutes after the injection. Symptoms may include:
 - trouble breathing
 - narrowing of airways
 - stomach cramps
 - sweating
 - numbness of your mouth
 - pain (e.g., back and chest)
 - feeling anxious
 - feeling warm
 - rash
 - feeling light-headed or feeling like you are going to pass out (faint)
 - blood pressure changes
- **Liver problems.** People with a history of hepatitis B or C virus or people who have certain liver function test changes may have an increased risk of developing new or worsening changes in certain liver tests during CABENUVA treatment. Liver problems have also happened in people without history of liver problems or other risk factors. Your healthcare provider may do blood tests to check your liver function. **Call your healthcare provider right away if you develop any of the following signs or symptoms of liver problems:**
 - your skin or the white part of your eyes turns yellow (jaundice)
 - dark or "tea-colored" urine
 - light-colored stools (bowel movements)
 - nausea or vomiting
 - loss of appetite
 - pain, aching, or tenderness on the right side of your stomach area
 - itching
- **Depression or mood changes.** Call your healthcare provider or get emergency medical help right away if you have any of the following symptoms:
 - feeling sad or hopeless
 - feeling anxious or restless
 - have thoughts of hurting yourself (suicide) or have tried to hurt yourself

The most common side effects of CABENUVA include:

- pain, tenderness, hardened mass or lump, swelling, redness, itching, bruising, and warmth at the injection site
- fever
- tiredness
- headache
- muscle or bone pain
- nausea
- sleep problems
- dizziness
- rash

These are not all the possible side effects of CABENUVA. Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

GET MORE INFORMATION

- Talk to your healthcare provider or pharmacist.
- Go to CABENUVA.com or call 1-877-844-8872, where you can also get FDA-approved labeling.

February 2022 CBN:1PIL

Trademark is owned by or licensed to the ViiV Healthcare group of companies.

©2022 ViiV Healthcare or licensor.

CBROGM210118 February 2022

Produced in USA.